


Natural Product Synthesis Through (4+3) Cycloaddition: (+/-)-Frondosin B and (+/-)-Liphagal

Duchan R. Laplace^a, Bart Verbraeken^a, Kristof Van Hecke^b
and Johan M. Winne^a

a. Department of Organic Chemistry, Laboratory for Organic Synthesis Ghent University, Krijgslaan 281 S4, 9000 Ghent, Belgium

b. Department of Inorganic Chemistry, Ghent University, Krijgslaan 281 S3, 9000 Gent, Belgium

Previous work in our laboratory revealed a new type of (4+3) cycloaddition that proceeds via an oxyallyl-type cation producing hard to synthesize 7-membered rings. This synthetic method is very useful in the field of natural product (NP) synthesis since many of these NPs contain furan fused cycloheptenes or straightforward synthetic derivatives thereof. The scope of this conversion has already been demonstrated using simple furanol precursors and diene systems.^[1]



A more recent study aimed at more elaborate polycycles containing cyclohepta[b]furan substructures in accordance to existing NPs. We envisaged a total synthesis for frondosin B and liphagal as obvious targets for our method, also the latter NP was identified as inhibitor for one isoform of the phosphatidylinositol 3-kinase (PI3K) enzyme and had already been the subject of research.^[2] Herein, we present a full account of our investigations towards a new and short synthesis of (+/-)-5epi-liphagal and frondosin B.^[3]

References

- [1] Winne, J. M., Catak, S., Waroquier, M. and Van Speybroeck, V. (2011), *Angew. Chem. Int. Ed.*, 50: 11990–11993.
- [2] Day, J. J., McFadden, R. M., Virgil, S. C., Kolding, H., Allewa, J. L. and Stoltz, B. M. (2011), *Angew. Chem. Int. Ed.*, 50: 6814–6818.
- [3] Laplace, D.R., Verbraeken, B., Vanhecke, V.H., Winne, J.M. (2014), *Chem. Eur. J.*, 20: 253-262.